

## REFERENCES CITED

- Abouheif, E. 1999. A method for testing the assumption of phylogenetic independence in comparative data. *Evolutionary Ecology Research*, 1: 895–909.
- Begon, M., Townsend, C. R. and Harper, J. L. 2006. *Ecology: from individuals to ecosystems*. Fourth edition. Blackwell Publishing.
- Bennett, A. F. 2003. *Linkages in the landscape: the role of corridors and connectivity in wildlife conservation*. Gland, Switzerland and Cambridge, UK: IUCN.
- Bock, C. E. and Ricklefs, R. E. 1983. Range size and local abundance of some North American songbirds: a positive correlation. *American Naturalist*, 122: 295–299.
- Böhning-Gaese, K., Caprano, T. and van Ewijk, K. 2006. Range size: disentangling current traits and phylogenetic and biogeographic factors. *American Naturalist* 167: 555–567.
- Brown, J. H. 1984. On the relationship between abundance and distribution of species. *American Naturalist* 124: 255–279.
- Burnham, K. P. and Anderson, D. R. 2002. *Model selection and multimodel inference: a practical information-theoretic approach*. Springer.
- Bush, M., Silman, M. R. and Urrego, D. H. 2004. 148,000 Years of Climate and Forest Change in a Biodiversity Hot Spot. *Science* 303: 827–829.
- Calendini, F. and Martin, J.-F. 2005. PaupUP v1.0.3.1 A free graphical frontend for Paup\* Dos software.
- CBOL Plant Working Group. 2009. A DNA barcode for land plants. *Proceedings of the National Academy of Sciences, USA*, 106: 12794–12797.
- Chase, J. M. and Leibold, M. A. 2003. *Ecological Niches: Linking Classical and Contemporary Approaches*. University of Chicago Press, IL.
- Chown, S. L. 1997. Speciation and rarity: separating cause from consequence. Pp. 91–109 in W. E. Kunin and K. J. Gaston (eds), *The Biology of Rarity: Causes and Consequences of Rare and Common Differences*. Chapman and Hall, London.
- Chown, S. L. and Gaston, K. J. 2000. Areas, cradles and museums: the latitudinal gradient in species diversity. *Trends in Ecology and Evolution* 15: 311–315.
- Clausing, G. and Renner, S. S. 2001. Molecular phylogenetics of Melastomataceae and Memecylaceae: Implications for character evolution. *American Journal of Botany* 88: 486–498.

Collingham, Y. C. and Huntley, B. 2000. Impacts of habitat fragmentation and patch size upon migration rates. *Ecological Applications*, 10: 131–144.

Collingham, Y. C., Hill, M. O. and Huntley, B. 1996. The migration of sessile organisms: a simulation model with measurable parameters. *Journal of Vegetation Science*, 7: 831–846.

Crawley, M. J. 2002. Statistical Computing: an Introduction to Data Analysis Using S Plus. Wiley-IEEE Press.

Davies, T. J., Purvis, A., and Gittleman, J. L. 2009. Quaternary climate change and the geographic ranges of mammals. *American Naturalist* 174: 297–307.

de Queiroz, K. 1998. The general lineage concept of species, species criteria and the process of speciation. Pp. 57–75 in D. J. Howard and S. H. Berlocher (eds.), *Endless forms*. Oxford University Press, NY.

de Queiroz, K. 2005. A unified concept of species and its consequences for the future of taxonomy. *Proceedings of the California Academy of Sciences* 56: 196–215.

de Queiroz, K. 2007. Species concepts and species delimitation. *Systematic Biology* 56: 879–86.

Draper, N. R. and Smith, H. 1998. *Applied Regression Analysis*. 1 John Wiley and Sons, Inc., NY.

Drummond, A. J. and Rambaut, A. 2007. BEAST: Bayesian evolutionary analysis by sampling trees. *BMC Evolutionary Biology* 7: 214, <http://beast.bio.ed.ac.uk/>.

Drummond, A. J., Ho, S. Y. W., Phillips, M. J. and Rambaut, A. 2006. Relaxed Phylogenetics and Dating with Confidence. *PLoS Biol* 4(5): e88. doi:10.1371/journal.pbio.0040088

Fazekas A. J., Burgess, K. S., Kesaniakurti, P. R., Graham, S. W., Newmaster, S. G., Husband, B. C., Percy, D. M., Hajibabaei, M. and Barrett, S. C. H. 2008. Multiple multilocus DNA barcodes from the plastid genome discriminate plant species equally well. *PLoS ONE* 3, e2802.

Felsenstein, J. 1985. Phylogenies and the comparative method. *American Naturalist* 125: 1–15.

Foote, M., Crampton, J. S., Beu, A. G., Marshall, B. A., Cooper, R. A., Maxwell, P. A. and Matcham, I. 2007. Rise and fall of species occupancy in Cenozoic fossil mollusks. *Science* 318: 1131–1134.

Fortin, M. J. and Dale, M. 2005. *Spatial analysis for ecologists*. Cambridge University Press, Cambridge.

Garland, T., Harvey, P. H. and Ives, A. R. 1992. Procedures for the analysis of comparative data using phylogenetically independent contrasts. *Systematic Biology*, 41: 18–32.

- Gaston, K. J. 1991 How large is a species' geographic range? *Oikos* 61: 434–438.
- Gaston, K. J. 1994. Measuring geographic range sizes. *Ecography* 17: 198–205.
- Gaston, K. J. 1998. Species-range size distributions: products of speciation, extinction and transformation. *Philosophical Transactions of the Royal Society B: Biological Sciences* 353: 219–230.
- Gaston, K. J. 2003. The structure and dynamics of geographic ranges. Oxford University Press, Oxford.
- Gaston, K. J. 2008. Biodiversity and extinction: the dynamics of geographic range size. *Progress in Physical Geography* 32: 678–683.
- Gaston, K. J. 2009. Geographic range limits: achieving synthesis. *Proceedings of the Royal Society B* 276: 1395–1406.
- Gaston, K. J., Blackburn, T. M., Greenwood, J. J. D., Gregory, R. D., Quinn, R. M. and Lawton, J. H. 2000. Abundance-occupancy relationships. *Journal of Applied Ecology* 37: 39–59.
- Gaston, K. J., Blackburn, T. M. and Lawton, J. H. 1997. Interspecific abundance-range size relationships: an appraisal of mechanisms. *Journal of Animal Ecology* 66: 579–601.
- Gaston, K. J. and Fuller, R. A. 2009. The size of species geographic ranges. *Journal of Applied Ecology* 46: 1–9.
- Grace, J. B. 2006. Structural equation modeling and natural systems. Cambridge University Press.
- Grafen, A. 1989. The phylogenetic regression. *Philosophical Transactions of the Royal Society of London B*, 326: 227–241.
- Hanski, I. 1982. Dynamics of regional distribution: the core and satellite species hypothesis. *Oikos*, 38: 210–221.
- Hanski, I., Kouki, J. and Halkka, A. 1993. Three explanations of the positive relationship between distribution and abundance of species. Pp. 108–116 in R. Ricklefs and D. Schlüter (eds.), *Historical and Geographical Determinants of Community Diversity*. University of Chicago Press, Chicago.
- Harvey, P. H. and Pagel, M. D. 1991. *The Comparative Method in Evolutionary Biology*. Oxford University Press, New York.
- Heino, J. 2005. Positive relationship between regional distribution and local abundance in stream insects: a consequence of niche breadth or niche position? *Ecography* 28: 345–354.

- Hijmans, R. J., Cameron, S. E., Parra, J. L., Jones, P. G. and Jarvis, A. J. 2005. Very high resolution interpolated climate surfaces for global land areas. International Journal of Climatology 25: 1965–1978.
- Hirzel, A., Hausser, J. and Perrin, N. 2001. Biomapper 1.0. Lausanne, laboratory for conservation biology, URL: <http://www.unil.ch/biomapper>.
- Hirzel, A.H., Hausser, J., Chesser, D. and Perrin, N. 2002. Ecological-niche factor analysis: how to compute habitat-suitability maps without absence data? Ecology 83: 2027–2036.
- Holt, R. D. and Keit, T. H. 2005. Species' borders: a unifying theme in ecology. Oikos 108: 3–6.
- Honnay, O., Bossuyt, B., Verheyen, K., Jacquemyn, H., Butaye, J. and Hermy, M. 2002. Ecological perspectives for the restoration of plant communities in European temperate forests. Biodiversity & Conservation 11: 213–242.
- Hooghiemstra, H., Wijninga, V. M. and Cleef, A. M. 2006. The paleobotanical record of Colombia: implications for biogeography and biodiversity. Annals of the Missouri Botanical Garden 93: 297–324.
- Hubbell, S. P. 2001. The unified neutral theory of biodiversity and biogeography. Princeton University Press, Princeton, NJ.
- Huelsenbeck J. P. and Ronquist, F. 2001 MRBAYES: Bayesian inference of phylogenetic trees. Bioinformatics 17: 754–755.
- Jablonski, D. 1987 Heritability at the species level: analysis of geographic ranges of Cretaceous mollusks. Science 238: 360–363.
- Jackson, S. T. and Overpeck, J. T. 2000. Responses of plant populations and communities to environmental changes of the late Quaternary. Paleobiology 26: 194–220.
- Jetz, W. and Rahbek, C. 2002: Geographic range size and determinants of avian species richness. Science 297: 1548–1551.
- Johnson, C. N. 1998. Species extinction and relationship between distribution and abundance. Nature 394: 272–274.
- Jones, K. E., Sechrest W. and Gittleman, J. L. 2005. Age and area revisited: identifying global patterns and implications for conservation. Pp. 142–165 in A. Purvis, J. L. Gittleman and T. Brooks (eds.), Phylogeny and Conservation. Cambridge University Press, Cambridge.
- Kreft, H., Sommer, J. H. and Barthlott, W. 2006: The significance of geographic range size for spatial diversity patterns in Neotropical palms. Ecography 29: 21–30.

Kress J. W., Wurdack, K. J., Zimmer, E. A., Weigt, L. A. and Janzen, D. H. 2005. Use of DNA barcodes to identify flowering plants. *Proceedings of the National Academy of Sciences, USA*, 102: 8369–8374.

Kress W. J. and Erickson, D. L. 2007. A two-locus global DNA barcode for land plants: The coding *rbcL* gene complements the non-coding *trnH-psbA* spacer region. *PLoS ONE* 2:e508.

Lennon, J. J., Koleff, P., Greenwood, J. J. D. and Gaston, K. J. 2004: Contribution of rarity and commonness to patterns of species richness. *Ecology Letters* 7: 81–87.

Liow, L. H. 2007. Does versatility as measured by geographic range, bathymetric range and morphological variability contribute to taxon longevity? *Global Ecology and Biogeography* 16: 117–128.

Liow, L. H. and Stenseth, N. C. 2007. The rise and fall of species: implications for macroevolutionary and macroecological studies. *Proceedings of the Royal Society B* 274: 2745–2752.

Lomolino, M. V., Riddle, B. R. and Brown, J. H. 2006. *Biogeography*. Sinauer Associates, Sunderland.

MacKenzie, D. I., Nichols, J. D. ,Royle, J. A., Pollock, K. H., Bailey, L. L. and Hines, J. E. 2006. Occupancy estimation and modeling: inferring patterns and dynamics of species occurrence. Elsevier, Amsterdam.

McKinney, M. L. 1997. How do rare species avoid extinction? A paleontological view. Pp 110–129 in W. E. Kunin and K. J. Gaston (eds.), *The biology of rarity: causes and consequences of rare common differences*. Chapman Hall, London.

Maddison, W. P. and Maddison, D. R. 2006. Mesquite: A modular system for evolutionary analysis. Version 1.1. <http://mesquiteproject.org>.

Margules, C. R. and Pressey, R. L. 2000. Systematic conservation planning. *Nature* 405: 243–253.

Martinez-Meyer, E., Peterson, A. T. and Hargrove, W. W. 2004. Ecological niches as stable distributional constraints on mammal species, with implications for Pleistocene extinctions and climate change projections for biodiversity. *Global Ecology and Biogeography* 13: 305–314.

Midford, P. E., Garland, T. and Maddison, W. P. 2005. PDAP Package of Mesquite. Version 1.07.

Murray, B. R. and Hose, G. C. 2005. Life-history and ecological correlates of decline and extinction in the endemic Australian frog fauna. *Austral Ecology* 30: 564–571.

- Pagel, M. D. 1992. A method for the analysis of comparative data. *Journal of Theoretical Biology*, 156: 431–442.
- Paul, J. R., Morton, C., Taylor, C. M. and Tonsor, S. J. 2009. Evolutionary time for dispersal limits the extent but not the occupancy of species' potential ranges in the tropical plant genus *Psychotria* (Rubiaceae). *American Naturalist* 173: 188–199.
- Paul, J. R. and Tonsor, S. J. 2008. Explaining geographic range size by species age: a test using Neotropical *Piper* species. Pp. 46–62 in W. P. Carson and S. Schnitzer (eds.), *Tropical forest community ecology*. Blackwell, London.
- Payne, J. L. and Finnegan, S. 2007: The effect of geographic range on extinction risk during background and mass extinction. *Proceedings of the National Academy of Sciences of the United States of America* 104: 10506–10511.
- Peterson, A. T., Papes, M. and Soberón, J. 2008. Rethinking receiver operating characteristic analysis applications in ecological niche modeling. *Ecological Modelling* 213: 63–72.
- Phillips, S., Anderson, J., R. P. and Schapire, R. E. 2006. Maximum entropy modelling of species geographic distributions. *Ecological Modelling*, 190: 231–259.
- Phillips, S. J. and Dudik, M. 2008. Modelling of species distributions with Maxent: new extensions and a comprehensive evaluation. *Ecography* 31: 161–175.
- Pulliam, H. R. 1988. On the relationship between niche and distribution. *Ecology Letters* 3: 349–361.
- Pohlman, C. L., Nicotra, A. B. and Murray, B. R. 2005: Geographic range size, seedling ecophysiology and phenotypic plasticity in Australian Acacia species. *Journal of Biogeography* 32: 341–51.
- Quinn, G. P. and Keough, M. J. 2002. *Experimental Design and Data Analysis for Biologists*. Cambridge University Press.
- R Development Core Team. 2006. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. <http://www.R-project.org>
- Reeve, J. and Abouheif, E. 2003. Phylogenetic independence. Version 2.0. Available from <http://biology.mcgill.ca/faculty/abouheif/> accessed August 2006.
- Renner, S. S. 2004. Bayesian analysis of combined data partitions, using multiple calibrations, supports recent arrival of Melastomataceae in Africa and Madagascar. *American Journal of Botany* 91: 1427–1435.

Renner, S. S., Clausing, G. and Meyer, K. 2001. Historical biogeography of Melastomataceae: the roles of tertiary migration and long-distance dispersal. *American Journal of Botany* 88: 1290–1300.

Renner, S. S. and Meyer, K. 2001. Melastomeae come full circle: biogeographic reconstruction and molecular clock dating. *Evolution* 55: 1315–1324.

Rheindt, F. E., Ulmar, G. T. and Abouheif, E. 2004. Rapidly evolving traits and the comparative method: how important is testing for phylogenetic signal? *Evolutionary Ecology Research* 6: 377–396.

Rosenzweig, M. L. 1995. Species diversity in space and time. Cambridge University Press, Cambridge.

Shaw, J., Lickey, E. B., Beck, J. T., Farmer, S. B., Liu, W., Miller, J., Siripun, K. C., Winder, C. T., Schilling, E. E. and Small, R. L. 2005. The tortoise and the hare II: Relative utility of 21 noncoding chloroplast DNA sequences for phylogenetic analysis. *American Journal of Botany* 92: 142–166.

Shugart, H. H. and Patten, B. C. 1972. Niche quantification and the concept of niche pattern. Systems analysis and simulation in ecology. Pp. 284–327 in B. C. Patten (ed.), Volume 2. Academic Press, New York.

Skov, F. and Svenning, J. C. 2004. Potential impact of climate change on the distribution of forests herbs in Europe. *Ecography* 27: 366–380.

Stamatakis, A., Hoover, P. and Rougemont, J. 2008. A fast bootstrapping algorithm for the RAxML Web-servers. *Systematic Biology* 57: 758–771.

Standards and Petitions Working Group. 2006. Guidelines for using the IUCN red list categories and criteria. Version 6.1. World Conservation Union SSC Biodiversity Assessments Subcommittee, Gland, Switzerland.

<http://app.iucn.org/webfiles/doc/SSC/RedList/RedListGuidelines.pdf>.

Stevens, G. C. 1989: The latitudinal gradient in geographical range: how so many species co-exist in the tropics. *American Naturalist* 133: 240–256.

Svenning, J.-C., Normand, S. and Skov, F. 2008. Postglacial dispersal limitation of widespread forest plant species in nemoral Europe. *Ecography* 31: 316–326.

Svenning, J.-C. and Skov, F. 2004. Limited filling of the potential range in European tree species. *Ecology Letters*, 7: 565.

Swofford, D. L. 2000. PAUP\*. Phylogenetic Analysis Using Parsimony (\*and Other Methods). Version 4. Sinauer Associates, Sunderland, Massachusetts.

Taberlet, P., Gielly, L., Patou, G. and Bouvet, J. 1991. Universal primers for amplification of three noncoding regions of chloroplast DNA. *Plant Molecular Biology*. 17: 1105-1109.

Tamura K, Dudley, J., Nei, M. and Kumar, S. 2007. MEGA4: Molecular Evolutionary Genetics Analysis (MEGA) software version 4.0. *Molecular Biology and Evolution* 24:1596-1599.  
Tropicos, Missouri Botanical Garden VAST Vascular Tropicos, Nomenclatural database. – <http://www.tropicos.org>.

Ulloa Ulloa, C. 2007. *Brachyotum sertulatum*, una especie nueva de Melastomataceae de Colombia. *Anales del Jardín Botánico de Madrid* 64: 69–73.

Vrba, E. S. and deGusta, D. 2004. Do species populations really start small? New perspectives from the Late Neogene fossil record of African mammals. *Philosophical Transactions of the Royal Society B* 359: 285–292.

Waldron, A. 2007: Null models of geographic range size evolution reaffirm its heritability. *American Naturalist* 170: 221–231.

Webb, T. J. and Gaston, K. J. 2000. Geographic range size and evolutionary age in birds. *Proceedings of the Royal Society B* 267: 1843–1850.

Webb, T. J. and Gaston, K. J. 2003. On the heritability of geographic range sizes. *American Naturalist* 161: 553–566.

Willis, J. C. 1922 Age and area: a study in geographic distribution and origin of species. Cambridge, UK: The University Press.

Wurdack, J. J. 1953. A revision of the genus *Brachyotum* (Tibouchineae - Melastomataceae). *Memoirs of the New York Botanical Garden* 8: 343–407.